

BACTERIOLOGICAL EVALUATIONS OF RESTRUCTURED FISH PRODUCT PROCESSED BY TUMBLING WITH EGG WHITE

YETIM H. and OCKERMAN H.W.

The Ohio State University, Department of Animal Science, Columbus, Ohio, USA

S-VIB.49

SUMMARY

Objectives were to develop an understanding of contamination levels and to identify organisms of concern in a restructured fish product which was tumbled with egg white. Fresh Channel Cat fish fillets were diced, and the proper amount of ingredients [2% salt, seasoning and 1% egg white (not in control group)] were mixed with the tissue before an intermittent (40 min rest, 20 min work) tumbling process (12 h at approximately 6°C.) Total aerobic plate count (APC), coliform, *Pseudomonas*, *Salmonella*, total psychrophilic, lipolytic and proteolytic bacteria were determined before and after tumbling. The tumbling process significantly influenced bacterial numbers when sampled before the cooking process and no *Salmonella* organism was detected in any sample. APC and *Pseudomonas* numbers were lower in the processed fish muscle when compared to the fresh control. Tumbling also caused an inhibition in APC, *Pseudomonas*, lipolytic and proteolytic growth during the process probably due to better cure distribution and/or faster cure penetration. There was a significant positive correlations between APC and *Pseudomonas*, and also between *Pseudomonas*, proteolytic and lipolytic bacteria numbers. In conclusion, the tumbling technique would be useful in improving bacteriological quality of an acceptable restructured fish product.

INTRODUCTION

It has been suggested that the fish industry should become much more innovative and produce an increasing number of interesting, value-added products for today's consumer. As new products are developed it is also the responsibility of the food processor to assure the consumer that they are safe and wholesome. Previous microbiological analysis results have indicated that microflora of seafood products varies considerably. This flora both quantitatively and qualitatively depends on the living environment and harvesting technique used for the seafood (Foster et al., 1977; Love, 1988) and on its storage conditions after processing.

The flesh of healthy, living fish is sterile, but spoilage bacteria are usually introduced during capture, handling or contact with equipment or people, and the microorganism become established rapidly (Shaw et al., 1984). Fish might be a carrier of many spoilage and water-borne pathogenic bacteria and their numbers and proliferation can generally be attributed to the following factors; (a) fish from polluted waters, (b) improper refrigeration, (c) improper sanitation procedures due to either facilities or handlers during the harvest or processing and (d) improper cooking. Obviously these factors effects not only shelf life but also microbial quality and wholesomeness of the product. Many research projects have been conducted to understand the microorganisms involved with public health and spoilage of seafood items, and these studies must continue to seek ways to mitigate the effects of contamination and natural flora through effective surveillance or control techniques (Shewan, 1976; Abeyta, 1983; Liston, 1990). This research was undertaken to develop an understanding of the level of contamination and the effects of processing on microbial numbers, and to identify some organisms of concern (spoilage and pathogen) in a fish product which was tumbled in the presence of egg white.

MATERIALS and METHODS

a. Sample Preparation: Fresh Channel Catfish fillets were first inspected for any dirt and bones, diced (approximately 3x3x2 cm), and then sampled for fresh analysis. The treatment groups were: 1) non tumbled and no egg white added (control); 2) no egg white added and tumbled; 3) non tumbled and egg white added; and 4) tumbled and egg white added. A proper amount of ingredients [2% salt, 1% egg white powder (in egg white batches), 1% sucrose, 0.5% white pepper, 0.1% garlic, 0.25% sodium tri-polyphosphate, 0.01% nitrite and 0.01% natural lemon flavor] were mixed with the product. The intermittent tumbling procedure (40 min rest, 20 min work) was conducted for 12h at 6°C on tumbled batches, and non tumbled batches were stored for the same time in the same place. Then, each of the treatments was immediately sampled for bacteriological analysis, and the product was stuffed into casing and cooked until an internal temperature of 77°C was attained. After cooling, the product was sampled again for thermophilic bacteria enumeration.

b. Bacteriological Analysis: American Public Health Association (APHA, 1992) procedures were used to identify and count the bacteria, other than *Salmonella*, investigated or monitored in this research. Serial dilutions of the samples in 0.1% peptone water were plated in duplicate and incubated according to the procedures shown in Table 1. For *Salmonella* determinations the BAM (Bacteriological Analysis Manual) procedure was followed (FDA, 1984). Twenty five grams of the sample was first enriched in Selenite Cystine broth, and then streaked on pre-poured agar (XLD and BSA) plates. The suspicious colonies were subjected to further biochemical evaluations for confirmation.

c. Experimental Procedure and Statistical Analysis: Fresh fish fillets and 4 different treatments (non-tumble and tumble fish muscle with or without egg white) were tested. The experiment consisted of a (1x2x2) factorial block design (fresh, tumbling and egg white) with five replications. The individual fish batches are considered as a block, and ANOVA techniques were applied to the logarithms (base 10) of the bacterial numbers. Differences in treatment means and correlations were determined using SAS (SAS, 1985). Duncan's multiple range test was used for multiple comparison of means (Duncan, 1955). Additionally, if no interaction term was significant, tumbling and egg white treatments data were combined where appropriate, and another statistical analysis was performed on the treatments to see the influence of these factors.

RESULTS AND DISCUSSION

As can be seen from Table 2, there was significant changes in bacteria numbers between the treatments and fresh sample. Total Aerobic bacteria (APC) number was higher in fresh tissue compared to the processed groups and it was lowered with the tumbling process. The reason for this reduction in the processed treatments might be the curing ingredients such as salt, nitrite, phosphate, sugar and garlic which may cause an inhibition on colony forming units (APHA, 1992). Total psychrophilic bacteria number continued to increase during processing probably due to processing temperature which was 6 °C, but the increasing rate was not very high in the tumbled treatments. Also, *Pseudomonas* bacteria numbers were low in the tumbling process samples when compared to the original (in fresh fish) number, and it was also lower than the non-tumbled counterparts. In general, a lower bacterial number was determined in the tumbled groups compared to the non-tumbled treatments. The reason for this outcome might be a faster cure distribution or penetration into the muscle pieces (Mills et al. 1980; Knipe et al. 1981; Leak et al, 1984). However, coliform, proteolytic and lipolytic bacteria numbers were higher in processed groups irrespective of the treatments although tumbling showed a slightly lower bacterial count. The reason for this result might be unexpected contamination during the processing either from equipment or ingredients used in this study. Thermophilic bacteria numbers were quite low and were not significantly different among the treatments. Also, none of the samples including fresh tissue was *Salmonella* positive in the treatments although there are many reports of *Salmonella* incidence in seafood and fish products (Taylor, 1988; Garret, 1988; Liston, 1990; Nickelson, 1992).

The results obtained from the other statistical analysis in which the data was pooled is shown in Figure 1 and 2. As can be followed from Figure 1, egg white had no effect on the alteration of the bacteria enumerated in this research. However, tumbling had a significant effect on retarding proliferation of the bacteria enumerated in this study. The significant differences were in APC, *Pseudomonas*, lipolytic and proteolytic bacteria numbers probably due to better distribution of the cure ingredients with the tumbling process.

CONCLUSIONS

The utilization of tumbling significantly altered the bacteriological properties of restructured fish product during processing. Tumbling decreased APC, *Pseudomonas*, proteolytic and lipolytic bacteria while egg white addition had no significant effect on the enumerated bacteria. There were significantly positive relations between *Pseudomonas* and proteolytic bacteria, and between total psychrophiles and *Pseudomonas* numbers. Also, there was significant positive correlations between APC and *Pseudomonas* bacteria counts. These results confirm that the dominant bacteria genus in seafood was *Pseudomonas* which agrees with previously reported research by Shewan, 1976; Hobbs, 1983; Ward and Baj, 1988. It could be concluded from the bacteriological results that tumbling would be a useful technique to improve the microbial quality of fish products during processing.

REFERENCES

- Abeysa, C. 1983. Bacteriological quality of fresh seafood products from Seattle retail markets. *J. Food Prot.* 46: 901-909.
- APHA, 1992. Compendium of methods for the microbiological examination of foods. 3rd Ed., Edited by Vanderzant, C. and Splittstoesser, D.F. American Public Health Association, Washington, DC.
- Duncan, D.B. 1955. Multiple Range and F Test. *Biometrics.* 11: 1.
- Food and Drug Administration, 1984. *Bacteriological Analytical Manual*, 6th Ed. Arlington, VA.
- Foster, J.F., Fowler, J.L. and Dacey, J. 1977. A microbial survey of various fresh and frozen seafood products. *J. Food Prot.* 40: 300-303.
- Garret, S.E. 1988. Microbiological standards, guidelines and specifications and inspection of seafood products. *Food Technol.* 42 (3): 90-93.
- Hobbs, G. 1983. Microbial spoilage of fish. "In: *Food Microbiology; Advances and Prospects*, Edited by Roberts, T.A. and Skinner, F.A. pp 217, Academic Press, New York.
- Knipe, C.L. Plimpton, R.F. and Ockerman, H.W. 1981. Effects of tumbling and tumbling temperature on total microbial plate counts and quality of boneless, cured hams. *J. Food Sci.* 46: 212-215.
- Leak, F.W. Kemp, J.D. Langlois, B.E. and Fox, J.D. 1984. Effects of tumbling and tumbling time on quality and microflora of dry-cured hams. *J. Food Sci.* 49: 695-698.
- Liston, J. 1990. Microbial hazards of seafood consumption. *Food Technol.* 44 (12): 56-62.
- Love, R.M. 1988. *The Food Fishes: Their intrinsic variations and practical implications*. Farrand Press, London.
- Mills, E.W., Plimpton, R.F. and Ockerman, H.W. 1980. Residual nitrite and total microbial plate counts of hams as influenced by tumbling and four ingoing nitrite levels. *J. Food Sci.* 45: 1297-1300.
- Nickelson, R. 1992. Bacteria of significance in the international trade of shrimp. *Scope, Technical Bull. of Silliker Labs.* 7: (2) 1-5, 7.
- Ockerman, H.W. 1985. *Quality Control of Muscle Tissue*. The OSU. Dept. of Anim. Sci., Columbus, OH.
- SAS. 1985. *SAS User's Guide: Statistics*", 5th. Ed. SAS Institute, Inc. Raleigh, NC.

- Shewan, J.M. 1976. The bacteriology of fresh and spoiling fish and the biochemical changes induced by bacterial action. Proc. of the Conf. on the Handling, Proc. and Mark. of the Tropical Fish. July, 5-9, 1976, London.
- Shaw, S.J. Bligh, E.G. and Woyewoda, A.D. 1984. Effects of delayed filleting on quality of cod flesh. J. of Food Sci. 49: 979-980.
- Taylor, S.L. 1988. Marine toxins of microbial origin. Food Technol. 42 (3): 94-98
- Ward, D.R. and Baj, N.J. 1988. Factors affecting microbiological quality of seafood. Food Technol. 42 (3): 85-89.