

FACTORS CONTRIBUTING TO FADING AND DISCOLORATION OF SLICED, PACKAGED SALAMI UNDER ILLUMINATION

Oddvin Sørheim^{1*}, Solveig Uglem², Per Lea¹, Ingrid Måge¹, Marit Kvalvåg Pettersen¹ and Hanne Larsen¹

¹Nofima AS, Osloveien 1, NO-1431 Ås, Norway

²Grilstad AS, Ranheimsvegen 129B, NO-7053 Ranheim, Norway

*Corresponding author email: oddvin.sorheim@nofima.no

Abstract - Sliced salami packaged in modified atmospheres is prone to discoloration caused by the combination of residual O₂ in the headspace and light exposure at display. In this study, factors contributing to discoloration of salami were examined. The consumption of detrimental residual O₂ was faster at 20 than 4 °C. Light-induced discoloration increased with increasing levels of initial residual O₂ in the packages, and was higher at 4 than 20 °C. Salami in packages without O₂ present had a slight fading under light at 20 °C. To avoid discoloration of sliced, packaged salami, the packages should be stored in darkness until nearly all residual O₂ is consumed by the product. Thereafter, the product can safely be displayed under light without risking discoloration that would cause consumer rejection and food waste.

Key Words – colour, dry cured sausage, light exposure, residual oxygen.

I. INTRODUCTION

Fermented and dried salami sausages are available in many parts of the world with different processing and packaging. For sliced packaged salami, a red colour is regarded as one of the main criteria for purchase. Discoloration of the product can cause rejection by consumers, food waste and economical loss. Photooxidation of the pigment nitrosylmyoglobin is due to the combined effects of oxygen and light [1]. Modified atmosphere packaging (MAP) has become common for sliced salami, usually with 100 % N₂ or N₂ with small fractions of CO₂. Crucial for stabilizing colour of cold cuts in MAP is keeping residual O₂ in the headspace at a low level [2] and using packaging materials with high O₂ barrier properties [3].

In Norway, shelf stable raw fermented salami with a water activity below 0.90 is sold at room temperature, or in chilled cabinets. The display of dry cured sausages at various temperatures can affect colour and other quality parameters of the products.

The aim of this study was two-fold: to evaluate the effects of various levels of residual O₂ on the colour of sliced, packaged salami under light display, and elucidate possible non-O₂ related light effects on colour changes of the product.

II. MATERIALS AND METHODS

The study was designed with two experiments using sliced, packaged salami, the first with different concentrations of residual O₂ and the second without O₂. The samples were displayed under fluorescent light at 4 and 20 °C.

Salami sausages from Grilstad (Ranheim, Norway) were used in the study. The sausages had a diameter of 75 mm and contained ca. 35 % fat.

The first packaging experiment was conducted in the pilot plant at Nofima. Sausages were sliced and packaged on a Multivac R145 thermoforming machine (Multivac, Wolfertschwenden, Germany). The base web and top film named Multipet 450 and Biaxer 65 XX (both Wipak, Nastola, Finland) had O₂ transmission rates (OTR) of 10 and 5 cm³/m²/day at 23 °C and 50 % RH. Gas to product ratio was ca. 2:1. The packages were initially stored for 14 days at 4 °C in darkness to allow for complete removal of residual O₂ from headspace by internal O₂ consumption by the product.

Shortly before illumination, various levels of air corresponding to 0.1 to 1.0 % O₂ were injected into the packages through septas. Packages with air were made by puncturing with a needle.

The second packaging experiment was performed at Grilstad with a thermoforming machine, using high barrier base web and top film with OTR's below 5 cm³/m²/day at 23 °C and 50 % RH. These packages were stored in darkness, first at 10 °C for 1.5 months, followed by 4 and 20 °C for another 4.5 months, before being exposed to light with no residual O₂ in the headspace.

The illumination for both packaging experiments was performed in the Nofima lab. The light intensity was ca. 1000 lux continuous at the salami surfaces at both display temperatures. At 4 °C, Natura de luxe L36W/76 (Osram, Munich, Germany) recommended for meat display illumination was used. At 20 °C, Auralight T5 Supreme HO 49W/830 (Auralight Int., Karlskrona, Sweden) common for ceilings of grocery stores was used. The light exposure lasted for 4 and 13 days for the first and second packaging experiments, respectively. 4-5 packages were used for each series and test conditions.

Analyses included measurement of residual O₂ in the headspace of packages by using a Checkmate 3 instrument (Dansensor, Ringsted, Denmark). CIE a* redness values on the salami surfaces were analysed non-destructively through the top films with a Minolta Chroma Meter CR-400 (Konica Minolta, Inc., Tokyo, Japan) with a 8 mm viewing port, 2° viewer angle and illuminant D₆₅ with 4 replicate measurements per sample.

Statistics was performed in MATLAB (R2014b, The Mathworks, Inc., Nattick, MA, USA). a* values were evaluated by a fixed-effects ANOVA model with the main effects temperature, O₂ level and days of storage.

III. RESULTS AND DISCUSSION

In the first experiment, the rate of O₂ consumption was studied with a fixed initial O₂ headspace concentration of 1.0 %, and packages were displayed at 4 and 20 °C for 4 days under light

(Fig. 1). The data show that the O₂ consumption was dependent on temperature, with much faster decline in O₂ concentrations at 20 compared to 4 °C. The bacteriological activity and requirement of O₂ from starter cultures in the sausages were probably higher at 20 °C. In agreement with our study, the O₂ consumption of dry cured sausages under long term dark storage was higher at 22 than 4 °C [4].




Figure 1. Reduction of residual O₂ in the headspace of illuminated packages with sliced salami initially filled with 99 % N₂ and 1 % O₂. Symbols: ● at 4 °C and ■ at 20 °C.

The effect of different initial O₂ concentrations between 0.1 and 1.0 % on a* redness values after 4 days of light display is demonstrated in Fig. 2. a* values declined with increasing level of residual O₂. Samples in air, simulating a full leakage, had much lower a* values. With exception of samples in 0.25 and 21 % O₂, the redness was temperature dependent, with significantly (p < 0.05) higher a* values at 20 than 4 °C, due to the faster removal of residual O₂ at this high temperature (Fig. 1). An additional control with 1.0 % O₂ was stored in darkness for 4 days, and had a* values of ca. 16 both at 4 and 20 °C, not different (p > 0.05) from illuminated samples in 0 % O₂.




Figure 2. a* values of sliced, packaged salami with 0 – 21 % initial O₂ and displayed under light at 4 and 20 °C for 4 days. Symbols: ■ at 4 °C (left bars) and ■ at 20 °C (right bars). Bars with different letters are significantly different ($p < 0.05$).

In the second experiment, packages with sliced salami without residual O₂ in the headspace were illuminated for 13 days, and a* values were recorded (Fig. 3). Samples at 4 °C did not differ between 0, 6 and 13 days display ($p > 0.05$). However, samples at 20 °C started at higher a* values and redness than at 4 °C ($p < 0.05$), and declined by 1.5 – 2 a* values to days 6 and 13 ($p < 0.05$). This fading or slight loss of redness is not likely explained by photooxidation processes, but could be due to changes in surfaces properties like reflectance and interaction with fat and the melting point of this substance. The fading phenomena is of a much lower magnitude than the photooxidation by residual O₂ (Fig. 2), and is not likely to cause consumer rejection. Still, the data suggest that a slight fading of salami can take place even without measurable O₂ present in the headspace.




Figure 3. a* values of sliced, packaged salami without residual O₂ and displayed under light at 4 and 20 °C for 13 days. Symbols: ● at 4 °C and ■ at 20 °C.

The results of this study demonstrates that too high levels of residual O₂ in packages of sliced packaged salami at the start of illumination can cause discoloration, which increased with increasing levels of O₂. These findings are in agreement with results of cooked cured ham, where 0.15 % residual O₂ was set as maximum acceptable level [2]. Display at an elevated temperature, like 20 °C, reduced the extent of discoloration in the present salami, because of the higher consumption of detrimental residual O₂ at this temperature. However, other undesirable quality changes like fat oxidations and incipient fat melting can be a concern at higher temperatures.

IV. CONCLUSION

Sliced, packaged salami should be stored in darkness until all or nearly all residual O₂ in the headspace is consumed by the product. Light-induced discoloration was less severe at 20 than 4 °C, because internal O₂ consumption by the product was faster at the highest temperature.

ACKNOWLEDGEMENTS

The Norwegian Levy on Agricultural Products and the Norwegian Research Funding for Agriculture and Food

Industry (both Oslo, Norway) are thanked for funding the study through the projects FoodPack no. NFR 225351 and REforReM no. NFR 244418, respectively. The technical assistance of Aud Espedal at Nofima is highly appreciated.

REFERENCES

1. Møller, J.K.S. & Skibsted, L.H. (2002). Nitric oxide and myoglobins. *Chemical Reviews* 102: 1167-1178.
2. Larsen, H., Westad, F., Sørheim, O. & Nilsen, L.H. (2006). Determination of critical oxygen level on colour stability during chill storage of sliced, pasteurised ham packaged in modified atmosphere. *Journal of Food Science* 71: 407-413.
3. Yen, J.R., Brown, R.B., Dick, R.L. & Acton, J.C. (1988). Oxygen transmission rate of packaging films and light exposure effects on the color stability of vacuum-packaged dry salami. *Journal of Food Science* 53: 1043-1046.
4. Scetar, M., Kovacic, E., Kurek, M. & Galic, K. (2013). Shelf life of packaged sliced fermented sausage under different temperature. *Meat Science* 93: 802-809.