

ANTIOXIDANT SUPPLEMENTATION IS INEFFECTIVE TO REDUCE THE FREQUENCY OF PSE-LIKE ZONES IN PORK HAMS

A. Vautier^{1*}, T. Lhommeau¹, A. Promeprat¹, T. Sayd, V. Santé-Lhoutellier²

¹: IFIP – Institut du Porc, La Motte au Vicomte, BP 35104, 35651 Le Rheu Cedex, France

²: INRA, UR370 Qualité des Produits Animaux, F-63122 Saint Genès-Champanelle, France

I. INTRODUCTION

The frequency of pork hams with PSE-like zones is a major issue for the processed meat sector in France, despite recent improvements in the slaughter techniques and systematic sorting on the ultimate pH value of hams. Halothane genotype, still widespread in its heterozygous form, may play a significant role in this issue [1]. On the other hand, a recent work described at a molecular level a possible link between oxidative stress and the specific development of PSE-like zone [2]. From this hypothesis, an experimental design was carried out at the IFIP Experimental Station to evaluate the effect of antioxidant supplementation in pig diet on the appearance of PSE-like zone and more generally on the overall ham quality before and after cooking process.

II. MATERIALS AND METHODS

A batch of 204 pigs, from Piétrain sires which are homozygous for the halothane mutation *nn*, were fattened at the IFIP Experimental Station. The animals were fed with either a basal diet (Control group) or a basal diet supplemented with vitamin E (300 mg/kg), vitamin C (300 mg/kg) and organic selenium (0.2 mg/kg) during the nine last weeks of fattening (Antioxidant group). The two groups were slaughtered simultaneously after a 24h fasting. The pH1 (30 min. *post mortem* (*pm*)) and ultimate pH (pH24 *ext.*, at 24h *pm*) were measured at the external side of the *Semimembranosus*. Prior to deboning, the meat color was measured at 24h *pm* on the *Gluteus Medius* with a CR-400 colorimeter (Konica Minolta, Japan). After deboning, structural defects were evaluated on hams according to the PSE-like zone IFIP scale [3], then drip loss sampling (EZ method) was carried out on the internal side of the *Semimembranosus*. The drip loss was measured after 48h of storage at 6°C. Ultimate pH was also measured at the internal side of the *Semimembranosus* (pH24 *int.*). Cooking yield was measured following industrial “jambon cuit supérieur” process without individual traceability. Two batches of 98 hams were processed (Control group vs Antioxidant group) and sliced. The “paste-like” slice defect frequency was evaluated on the final product [1]. The effect of antioxidant was performed on meat quality traits (pH1, pH24 *int.*, pH24 *ext.*, L*a*b*) and on process traits (cooking yield, slicing defect rate) using the GLM procedure (SAS Institute, USA). The FREQ procedure was applied on PSE-like zone class data.

III. RESULTS AND DISCUSSION

We observed 5% of PSE meat (pH1<6.0) which is higher than the frequency generally observed (1.5% in a survey in 15 slaughterhouses in France [4]). The pH24 (*ext.*) is also lower than the average value observed on 1.6 million carcasses in 2015 (pH24=5.71 [5]). Considering these meat quality traits and the fact that all pigs were heterozygous for halothane genotype (*Nn*), it was not surprising to observe a total of 46% of hams showing severe PSE-like zone defects (class 3+4).

Table 1: Meat quality of the whole data set (n=204) and by PSE-like zone class – *Semimembranosus* muscle

	n=	m (sd)	PSE-like zone class ¹				p
			1	2	3	4	
pH1 (<i>ext.</i>)	203	6.27 (0.20)	6.34	6.26	6.28	6.17	0.02
pH24 (<i>ext.</i>)	194	5.64 (0.10)	5.74	5.66	5.60	5.56	<0.0001
pH24 (<i>int.</i>)	194	5.70 (0.17)	5.93	5.73	5.61	5.59	<0.0001
L* (<i>int.</i>)	193	58.8 (5.0)	52.3	56.8	62.0	64.3	<0.0001
Drip loss (%)	195	5.9 (2.0)	3.69	5.68	6.36	7.81	<0.0001

1: class 1=no defect; class 2=slight defect; class 3= deep defect on *Semimembranosus*; class 4= deep defect on several muscles (incl. *SM*)

PSE-like zones showed typical meat quality changes in comparison with meat free of defect: L* increase, and pH1, pH24, drip loss decrease (table 1). The dietary treatment did not affect pH24, L* and drip loss, in agreement with Monziols *et al.* [6] study focusing on vitamin E and selenium supplementation. However, pH1 differed significantly with a lower value in the Antioxidant group, for which no explanation was found. Our results were not in agreement with the study of Mourot *et al.* [7] who reported higher pH and lower drip loss with vitamin C supplementation.

Table 2: effect of antioxidant supplementation on meat quality of the *Semimembranosus* muscle

	Control	Antioxidant	p
pH1 (<i>ext.</i>)	6.30	6.23	0.01
pH24 (<i>ext.</i>)	5.65	5.62	0.06
pH24 (<i>int.</i>)	5.70	5.71	ns
L* (<i>int.</i>)	58.8	59.0	ns
Drip loss (%)	5.67	6.17	0.09

The antioxidant supplementation did not reduce the PSE-like zone frequency, therefore not confirming our hypothesis on a possible effect of antioxidant on the defect appearance. Despite a lower pH1 in Antioxidant group, a higher cooking yield was reported suggesting a protective effect of the diet on proteins (table 3). However, as reported on raw meat, the “paste-like” slice frequency was unchanged.

Table 3: effect of antioxidant supplementation on the frequency of PSE-like zones and processing yields

Frequency of PSE-like zone class (%)	Control	Antioxidant	p
1	15.8	12.1	
2	43.1	37.4	ns
3	33.7	33.3	
4	7.4	17.2	
Cooking yield (%)	84.7	85.4	0.05
“Paste-like” slice frequency (%)	40.2	35.9	ns

IV. CONCLUSION

In this study, the supplementation of antioxidant in the pig diet (vitamin E, vitamin C and organic selenium) did not reduce the frequency of PSE-like zone. However, the frequency of PSE-like zone was particularly high in this study (46%). It has been shown an additive effect of low pH24, low pH1 and halothane mutation status, being considered as major risk factors for PSE-like development.

REFERENCES

1. Vautier A., Gault E., Lhommeau T., Venduvre JL., Minvielle B. 2012. Impact of chilling rate and halothane genotype on the frequency of pse-like zones and the processing yields of hams. 58th ICoMST, Montreal, Canada
2. Théron L., Sayd T., Chambon C., Vénien A., Viala D., Astruc T., Vautier A., Santé-Lhoutellier V. 2018. Deciphering PSE-like muscle defect in cooked hams: a signature from the tissue to the molecular scale. Food Chemistry, submitted.
3. IFIP. 2005. Grille de notation du défaut « déstructuré » des muscles de la cuisse de porc.
4. Frotin P., Gault E., Chevillon P., Eon Y. 2007. Mise à jour des références techniques et des recommandations sur le process d'abattage dans 15 abattoirs. Rapport d'étude, 83p.
5. Vautier A. 2017. Vers un observatoire de la qualité des viandes françaises ? Journée Qualité de Viande IFIP, Rennes, France.
6. Monziols M., Quiniou N., Lebas N., Granier R., Vautier A. 2015. Effets d'une complémentation en sélénium et en vitamine E sur les performances de croissance et la qualité de la viande de porc. Journées de la Recherche Porcine.
7. Mourot J., Peiniau P., Aumaitre A. Chevillon P. 1992. Effet de l'apport de vitamine C sur les performances de croissance et la qualité de viande chez des porcs Large-White et croisés Large-White Piétrain. Journées de la Recherche Porcine.